



FGF10 gene

fibroblast growth factor 10

Normal Function

The *FGF10* gene provides instructions for making a protein called fibroblast growth factor 10 (FGF10). This protein is part of a family of proteins called fibroblast growth factors that are involved in important processes such as cell division, regulation of cell growth and maturation, formation of blood vessels, wound healing, and development before birth. By attaching to another protein known as a receptor, the FGF10 protein triggers a cascade of chemical reactions inside the cell that signals the cell to undergo certain changes, such as maturing to take on specialized functions. During development before birth, the signals triggered by the FGF10 protein appear to stimulate cells to form the structures that make up the ears, skeleton, organs, and glands in the eyes and mouth.

Health Conditions Related to Genetic Changes

lacrimo-auriculo-dento-digital syndrome

At least three mutations in the *FGF10* gene have been found to cause lacrimo-auriculo-dento-digital (LADD) syndrome. This disorder affects the formation of the lacrimal system (the system in the eyes that produces and secretes tears), the ears, the salivary glands (the glands in the mouth that produce saliva), the teeth, the hands, and sometimes, other parts of the body. The main features of LADD syndrome are abnormal tear production, malformed ears with hearing loss, decreased saliva production, small teeth, and hand deformities.

The *FGF10* gene mutations that cause LADD syndrome reduce the amount and activity of FGF10 protein. Less growth factor is available to bind to receptors, which decreases signaling within cells. A decrease in cell signaling disrupts cell maturation and development, which results in abnormal formation of the ears, skeleton, and glands in the eyes and mouth in people with LADD syndrome.

other disorders


Mutations in the *FGF10* gene have also been found to cause a condition similar to LADD syndrome called aplasia of lacrimal and salivary glands (ALSG). Individuals with this condition have absent (aplastic) or small (hypoplastic) lacrimal glands, which secrete tears in the eyes, and aplastic or hypoplastic salivary glands. This condition often causes eye irritability, chronic tearing (epiphora), dry mouth (xerostomia), and a greater susceptibility to cavities. Sometimes, other structures that are involved

in tear production are affected. The mutations that cause ALSG are thought to be less disruptive to cell signaling than the *FGF10* gene mutations that cause LADD syndrome, although in rare cases, the same mutation can cause either condition. Because there is overlap between the features of ALSG and LADD syndrome and because they are caused by mutations in the same gene, it is difficult to discern whether they are distinct disorders or part of the same disease spectrum.

Chromosomal Location

Cytogenetic Location: 5p12, which is the short (p) arm of chromosome 5 at position 12

Molecular Location: base pairs 44,301,655 to 44,389,780 on chromosome 5 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- FGF-10
- FGF10_HUMAN
- keratinocyte growth factor 2

Additional Information & Resources

Educational Resources

- Madame Curie Bioscience Database: Fibroblast Growth Factors
<https://www.ncbi.nlm.nih.gov/books/NBK6330/>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28FGF10%5BTIAB%5D%29+OR+%28fibroblast+growth+factor+10%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

OMIM

- APLASIA OF LACRIMAL AND SALIVARY GLANDS
<http://omim.org/entry/180920>
- FIBROBLAST GROWTH FACTOR 10
<http://omim.org/entry/602115>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_FGF10.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=FGF10%5Bgene%5D>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=3666
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/2255>
- UniProt
<http://www.uniprot.org/uniprot/O15520>

Sources for This Summary

- OMIM: FIBROBLAST GROWTH FACTOR 10
<http://omim.org/entry/602115>
- Milunsky JM, Zhao G, Maher TA, Colby R, Everman DB. LADD syndrome is caused by FGF10 mutations. Clin Genet. 2006 Apr;69(4):349-54.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16630169>
- Rohmann E, Brunner HG, Kayserili H, Uyguner O, Nürnberg G, Lew ED, Dobbie A, Eswarakumar VP, Uzumcu A, Ulubil-Emeroglu M, Leroy JG, Li Y, Becker C, Lehnerdt K, Cremers CW, Yüksel-Apak M, Nürnberg P, Kubisch C, Schlessinger J, van Bokhoven H, Wollnik B. Mutations in different components of FGF signaling in LADD syndrome. Nat Genet. 2006 Apr;38(4):414-7. Epub 2006 Feb 26. Erratum in: Nat Genet. 2006 Apr;38(4):495. Kubisch, Chriütian [corrected to Kubisch, Christian].
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16501574>
- Shams I, Rohmann E, Eswarakumar VP, Lew ED, Yuzawa S, Wollnik B, Schlessinger J, Lax I. Lacrimo-auriculo-dento-digital syndrome is caused by reduced activity of the fibroblast growth factor 10 (FGF10)-FGF receptor 2 signaling pathway. Mol Cell Biol. 2007 Oct;27(19):6903-12. Epub 2007 Aug 6.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17682060>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2099236/>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/FGF10>

Reviewed: June 2013
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services